

**Vinasse treatment in Brazil from 1970s to 1990s:
a lock-in case study within the ethanol agroindustry**

Abstract

Adding market value to a former by-product or waste must be an outcome of what is now being called an eco-innovation. Vinasse is an important by-product of the sugarcane processing in the manufacture of ethanol in Brazil. Until the 1970s, the main destinations of vinasse were surface waters and “sacrifice areas”. The fabulous augmentation in production of this effluent after the ProAlcool and the legal prohibition of vinasse discharges into rivers and other surface waters induced the exploration of new technological possibilities to its destination. During the two subsequent decades, several technological possibilities were under development. In the middle 80s, fertigation was held as the most diffused alternative and the problem was considered solved. This paper exposes the results of an enquiry into the technological trajectory building process that was guided by public and private decision making focused on the solution of the problem. Three main aspects were analyzed: the characterization of the vinasse destination problem in its various dimensions; the description of the relevant selection environment to the determination of the winner trajectory; and the identifications of the lock-in in the fertigation. It is recognized how other technological alternatives, mainly anaerobic biodigestion, representing the biotechnological via that had inspired many hopes, were abandoned. Besides, the risks of soil salinisation and groundwater aquifers contamination by continued fertigation could not be overlooked and offer evidence that the abandonment of other alternatives may have occurred too soon.

Keywords

Ethanol, vinasse, environmental impacts, scientific controversies, environmental regulation, *lock-in*

1. Introduction

Until the 1970s, the main destinations of vinasse or stillage (a by-product of ethanol production) were surface water and the so-called "sacrifice areas". With the extraordinary increase in production of the waste, after the implementation of the Pro-Alcohol, and with the statutory prohibition on release of vinasse in rivers in the late 70s, efforts were made to develop technological opportunities for its destination.

Over two decades, many technological possibilities were the subject of research and development. During the 1980s, a promising path was the development of the biodigestion, a biotechnological treatment of the waste that produces biogas.

Biogas produced by this biotechnological via failed to succeed in the market place. The mainstream literature concerned to examine the elements that hold back the creation and diffusion of cleaner technologies includes top-down and bottom-up arguments (Carrillo-Hermosilla *et al*, 2009). Top-down arguments are mainly developed by economists using aggregate economic models and bottom-up arguments are put forward by engineering studies. These arguments are considered too simplistic in focusing the dynamic of energy substitution and the technological change.

An alternative, evolutionary perspective has been proposed to overcome the shortcomings of the precedent approaches. This perspective explains the difficulties of creation and diffusion of cleaner technologies on the basis of the fact that the economic system can be locked-in to technological standards that are potentially environmentally inferior (Carrillo-Hermosilla *et al*, 2009).

This perspective, which was pioneered by David (1985) and Arthur (1988), is been developed by a number of lock-in studies, in a broad-understood evolutionary approach, mostly in the energy sector (Cowan & Kline, 1996; Jacobson & Johnson, 2000; Unruh, 2000; Jacobson & Bergek, 2004; Foxon *et al*, 2005; Raven, 2005; Van den Bergh *et al*, 2006;

Verbong & Geels, 2006) and in the transport sector (Schot *et al*, 1994; Cowan & Hulten, 1996; Hoogma *et al*, 2002).

Generally speaking, in the view of this literature, lock-in phenomena are due to the existence of significant increasing returns to adoption of existing technologies, produced by economies of scale, learning and networks. These economies must be understood by the integration of these technologies into the techno-economic system.

In this paper, we undertook a study of the formation of technological trajectory from public and private decision-making actions taken to solve the problem of disposal of vinasse and the development of new technologies.

For this study, it's useful to keep in mind the achievements which reveal that cleaner technologies face serious obstacles raised by the existence of increasingly high uptake performance due to the effects of economies of scale, learning and network externalities (Arthur, 1988; Kemp *et al*, 1998; Dolfsma & Leydesdorff, 2009). These obstacles must be understood by considering the co-evolution of the change processes, both technological and institutional (Carrillo-Hermosilla *et al*, 2009: 30-31).

More to the point, conventional economic approaches are being criticized in the view of the fact that the main questions are not optimization and equilibrium but path-dependent technological change within a context of co-evolving environmental, social and economic processes characterized both by irreversibility and uncertainty (Llerena & Matt, 1999; Mulder & Van Den Bergh, 2001; Frenken *et al*, 2004).

As has been shown by Montalvo (2007), the factors that hinder clean technologies go besides economic ones, including government policy, markets, communities and social pressure, attitudes and social values, technological opportunities and technological capabilities and organizational capabilities. It is been accepted that these factors, which can spur or hold back cleaner technologies, are intertwined as they may affect each other.

We argue that considering the co-evolution of the technological and institutional processes in the formation of the technological trajectory study requires a historical perspective and the consideration of the constitution of the relevant selection environment. Kemp *et al* (1998:177) note that selection environment is shaped by old technologies, through the emergence of production routines, existence of infrastructures, the formation of skills and habits, and established consumption patterns.

Along with the recognition of these views, we put forward that the constitution of the relevant selection environment keeps close connections with the scientific understanding of the environmental problem in cause and with the social pressures to solve it.

These are the fundamental reasons raised by the literature for which we particularly discuss here three aspects: the characterization of the problem of vinasse (in its various dimensions: historical, economic, environmental, scientific and social), the relevant selection environment in determining the winning technological trajectory, and the lock-in in fertigation.

The paper is organized into three sections. In the first one, it is carried out an extensive characterization of the problem of vinasse disposal, taking into account its distinct aspects: historical, economic, physical or environmental, social and scientific. The second section presents the analysis of the process of finding new technology solutions, focusing more attention on two competing technological paths: the fertigation and the anaerobic digestion. The third section presents the development of the selection environment which contributed to the establishment of the winning solution. The final discussion identifies the elements that have contributed to the stability of the existing option (lock-in) and which constitute obstacles to the success of the alternative option.

2. Characterizing the vinasse disposal problem

In this section, we characterize the environmental problem of the vinasse disposal in its distinctive aspects: historical-economic, physical or environmental and social and scientific ones.

2.1 Historical and economic aspects of vinasse disposal

Vinasse – or stillage – is a byproduct of the process of the ethanol production from the distillation of fermented juice of cane sugar.


For decades, even while it was not generated in the current volumes, vinasse was a cause of some health and environmental concern in the environmental control agencies and the scientific community in Brazil. The work of Almeida, 1952 (apud Szmrecsányi 1994), had already shown that the issue aroused the attention of scientists and was the object of studies in the 40s and 50s, the time when the waste was dumped into surface waters (a practice which been used for a long time) and "sacrifice areas."

With the implementation of the National Alcohol Program (ProAlcool), however, the environmental damage to flora and fauna from surface waters acquired an alarming dimension.

The National Alcohol Program was created, as we know, in order to promote the partial replacement of the gasoline used in light vehicles by hydrated alcohol as part of the actions taken by the Federal Government to reduce the impact of the rising oil prices in the 1970s.

Since the mid-70s until the late '80s, the spur to the ethanol production gave new impetus to the sugar cane agribusiness in the country. Figure 1 shows that with the implementation of the Pro-Alcohol, the scale of domestic production of ethanol has raised significantly. From the 638 million gallons per annum, which were produced on average in

the first half of the 70s, production increased dramatically at the end of the 90s, with the approximate volume of 16 billion gallons annually. Out of that amount about 15 billion is of ethanol fuel (Scheleder 1998).




Source: Corazza (2001).

The rise of the ethanol production had allowed the national production of alcohol cars reached 96% in 1985. However, the growing production of ethanol in Brazil has led inevitably to increased production of vinasse, worsening the problem of the waste disposal. As each liter of alcohol generates about 12 liters of vinasse, the growth of production of this waste was dramatic. The volume of vinasse generated annually in the country can be estimated, according to data presented by Hassuda (1989) in view of the production of alcohol at the end of the 90s, something around 192 billion gallons. The data on the path of the generation of vinasse are shown in Figure 2.

It should be noted that while there has been technological improvements in the sugarcane agro-industry since the implementation of ProAlcool, the ratio between the amount of vinasse generated per liter of alcohol showed no significant improvement, except in some plants, making them not only producers of alcohol (and sugar) but also of huge quantities of waste. Probably the fact that the sugar industry had not made significant efforts directed

towards reducing the proportion of alcohol and vinasse must be linked, on the one hand, to their heterogeneous nature and, secondly, to the lack of a more restrictive environmental regulations.

Figure 2 – Brazilian ethanol and vinasse production trends from 1967 a 1997. Averages were calculated in million of liters for harvests grouped each five years.



Source: Corazza (2001).

Interestingly, back in the 80's when National Alcohol Program was under a crisis, the environmental issue has tended to become an argument in favor of the program, such as was the case of the national energy independence issue in the early years of its implementation. Authors such as Cortez & Larson (1998), describing the environmental impacts of ethanol production, were enthusiastic about its advantages: the positive balance of CO₂, the power generation from a renewable source and the possibility of producing electricity through cogeneration plants.

For reasons like these, ProAlcool was praised by the World Commission on Environment and Development, having been quoted in the book "Our Common Future" as an example of exploitation of a renewable energy source.

If ProAlcool finds reasons to favor its continuity on environmental grounds listed by many authors, there are also those not so happy with other environmental impacts of the Program. Many authors, such as La Rovere (1981), Margulis (1982), Guarnieri & Januzzi (1992) and Szmrecsányi (1994), indicated negative impacts of the Program on environmental quality, particularly in regard to vinasse disposal.

2.2 Physical and environmental aspects of vinasse and its disposal

Vinasse is an effluent rich in organic matter and potassium, with significant levels of calcium, magnesium and sulfur and other minerals in small quantities. This is, in accordance with Plaza-Pinto (1999), a substance of suspended organic solids and minerals with high COD (Chemical Oxygen Demand), BOD (Biological oxygen demand), which explains its great pollution potential.

Along with pollution from organic substances, other physical aspects of vinasse are its low pH and the high temperature in which it is generated, resulting in its character of chemical pollutant (expressed by a high corrosion) and pollution of the physical nature (in the form of heat).

When it comes to the origin of the effluent, the sources are identifiable because they correspond to plants producing ethanol in the sugarcane agro-industry. As for the use of vinasse, the main sources are in sugarcane production, corresponding to multiple properties of planters. There is no survey on the amount of vinasse used in these properties annually, making it difficult to accurately identify those sources. In Sao Paulo, for example, the axis Campinas - Ribeirão Preto concentrates most of these properties.

The elements of the ecosystem that are potentially affected are surface water, soil and groundwater.

By the end of 70 years, increasing volumes of vinasse were released into surface waters. The proliferation of microorganisms due to exhaust the oxygen dissolved in water, destroying aquatic flora and fauna and hampers the supply of drinking water. Besides the stench, Almeida (*apud* Szmrecsányi, 1994) call attention to the worsening of endemic diseases such as malaria, amoebiasis and schistosomiasis.

The release of vinasse in surface waters were seasonal, following the cycle of ethanol production, which explains the acute nature of the problem, affecting the functions of self-regulation and self-reproduction of ecosystems. After the shock, along with the dilution of pollutants, fish populations seem to recover.

Another alternative for disposal of the vinasse is the so-called "sacrifice areas", surface areas that receive or are flooded with the untreated vinasse. Evidently, the "sacrifice areas" become completely useless for any other purposes.

Later, particularly from the 80s, with the spread of the practice of fertigation, which is going to be detailed later in this paper, and with the still continued use of "sacrifice areas", soil is the element of ecosystem that becomes the more affected by vinasse disposal.

Finally, we discuss, as we shall see, the possibility of contamination of groundwater sources. Given the controversial nature of this discussion, it will appear below.

2.3 Socio-scientific controversies of the vinasse disposal

While the effects of the discharge of vinasse on the water surface are well known to the point to raise no more disputes, the same does not seem to occur with the environmental impacts of their disposal in the soil or with the possibility of contamination of groundwater.

There are a number of scientific papers devoted to studying the effects of vinasse as fertilizer, while few research initiatives are turning to the effects of waste on groundwater quality.

We have observed that, according to studies of Gloria 1984 (*quoted* by Plaza-Pinto, 1999), the failure to observe the optimal doses of vinasse (which vary according to soil type and second sugarcane varieties), leads to risk of salinisation and degradation of the very quality of sugarcane produced. Despite this and although the wide diffusion of fertigation, Copersucar released in 1986, according to Plaza-Pinto (1999), the information that about 40% of the vinasse produced in the State of Sao Paulo is not yet exploited, and discarded in sacrifice areas, a practice that has been authorized with restrictions by the Company of Technology and Environmental Sanitation (Cetesb), which requires the use of waterproof blankets of PVC to cover the soil of these areas, for the protection of groundwater and surface waters. Given that the alcohol industry is a factor of vulnerability of aquifers Paulo (Hirata *et al*, 1991), this high percentage may be considered critical. Hirata *et al* (1991) show that the region that concentrates most of the sugar and alcohol in the State of São Paulo (axis Campinas - Ribeirão Preto) is located exactly on the recharge area of the Guarani Aquifer, one of the most important in South America. In this area, the groundwater is relatively close to the surface, which makes them vulnerable to infiltration of pollutants, especially the salts (potassium, nitrates, etc.). For this reason, Hassuda (1989), Cruz (1991), Righetto *et al* (1991) and Gloeden *et al* (1992) developed methods for monitoring the risk of groundwater contamination by leakage of vinasse and conducted studies on the possibility of contamination of groundwater aquifer belonging to the so-called Botucatu Aquifer (the region of the Guarani aquifer recharge). Although any contamination was found at the time, the results of these studies indicate leaching of nutrients from the vinasse into the groundwater, especially nitrates. For this reason, the authors reiterated the danger of further degradation of aquifers and were unanimous in stressing the need for more research on the risks of infiltration of vinasse in soil / fertigation for groundwater quality and human health.

It is appropriate to note the existence of the argument that the very cultivation of cane sugar would be responsible for absorbing very quickly most of the salts (especially potassium), and therefore prevent both the risk of salinity as well that of contamination of groundwater. However, given the total absence of control over the fertigation, it is not possible to confirm the validity of this argument.

Thus, there is controversy about soil salinisation and contamination of underground aquifers. The possible causal chains between the use of vinasse in soil, and the size, scope and timing of the appearance of adverse effects remain unclear.

It appears that the frequency and intensity of the release of vinasse in soil can influence the possibility that salts (potassium, nitrates, etc.) accumulate, producing, eventually, the two effects: salinisation and contamination of groundwater. This argument is, however, more research and scientific discussion, it seems, was doomed - at least so far - to oblivion.

3. Technological developments in the 1970s and 1980s

As pointed out by Margulis (1982), ProAlcool was not preceded by a stage of research with specific and coordinated programs in order to optimize production processes or to minimize waste and materials recovery. It was only after the establishment of its ambitious targets for increasing the production of alcohol (and vinasse), that several research centers have turned to the study of the problem of vinasse disposal. Thus, only when the immense amount of the waste (now being generated with the expansion of ethanol production after ProAlcool) showed the extent of the problem of its disposal, that efforts have been mobilized to develop technological possibilities to its solution. These possibilities included: aerobiosis, recycling of effluent in the fermentation process, the fertigation, the combustion, the

production of yeast, the use in construction, the manufacture of animal feed and anaerobic digestion.

Aerobiosis, recycling in fermentation and fertigation are simpler solutions from the technological standpoint, and sugarcane agro-industry had already some knowledge of them. These alternatives benefited also of some diffusion. It is likely that the low technological sophistication of these processes has made possible its use in industrial scale within the first years of ProAlcool.

Combustion, the production of yeast, the use of vinasse in civil construction and in the manufacture of animal feed, as well as the anaerobic digestion, were under differentiated stages of development, each in different degrees of maturity and intensity of research. It is possible that technological sophistication of these solutions had represented an obstacle to its dissemination within a sector traditionally of low technological dynamics.

In the next section, we discuss the main advantages and problem areas with which each of these alternatives faced, during their specific technological development efforts, which are briefly presented. Further, we describe and analyze the two alternatives that have gained more strength: the fertigation in terms of dissemination of its adoption, and anaerobic digestion, because it has mobilized expertise and institutions and provided a "bet" on the emergence of biotechnology as a technological paradigm.

The study of such technologies allows their classification into four groups according to the type of the waste treatment needed: 1) technologies that require concentration of the residue by evaporation, 2) aerobic fermentation, 3) anaerobic digestion, and 4) other uses for untreated vinasse.

The creation of new skills and mobilize different institutions behind these tracks or technological trajectories allows us to speak, to use the words of Dosi (1982), of a period of extraordinary search, or a pre-paradigmatic phase. We are able to speak also of the emergence

of capabilities in the technological domain as well as in the organizational field, two fundamental factors pointed by Montalvo (2007) affecting cleaner technologies.

4. The relevant selection environment definition

We feature here the relevant selection environment to the formation of the technological trajectory to solve the problem of disposal of vinasse.

The selection environment has to do with Carrillo-Hermosilla *et al* (2009) institutional environment that includes public policies, laws and organizations, which are taken into account here. But we go further by briefly characterizing: a) the technological aspects of the industry structure; b) a key macroeconomic element affecting the formation of the technological trajectory, as to say, oil prices, since petroleum is the main substitute energy source for ethanol; and c) a specific aspect of the institutional environment which consists of the level of institutionalization of the biotechnology at the time.

As we have already pointed out in this paper, the constitution of the relevant selection environment keeps close connections with the scientific understanding of the environmental problem in cause and with the social pressures to solve it, aspects developed in the previous sections. Here we try to shed some light in the connections of these aspects and the constitution of the selection environment.

In the alcohol sector, the agricultural and industrial phases do not always correspond to the same unit of venture. Thus, with regard to market structure, it is interesting to distinguish between two types of structure elements: the first is linked to units of origin of the residue, while the second is related to units of vinasse use. The first type of market structure involves the distilleries (which may be autonomous, while only producing alcohol, or attached, when they produce sugar and alcohol). The second type involves the planters and

therefore users of fertigation. Taking into account this distinction, it is possible to say that the industry has become more concentrated in the first and more sprayed in the second type.

With regard to the distilleries, we point out the technological duality between autonomous and attached distilleries. Different manufacturing processes make vinasse produced by independent (autonomous) distilleries more concentrated in pollutant load than that produced in attached distilleries: that is, the production of ethanol directly from the fermentation (in autonomous plants) generate more vinasse than indirect production from the fermentation of diluted molasses (in attached plants). With ProAlcool, since 1975 most of the plants was conceived as "autonomous", because it was not aimed to produce sugar, but ethanol. Consequently, most ethanol plants operating in the country are autonomous - that is, the type in which there is a larger ratio vinasse/ethanol. What is important to point out, however, is that for the industry as a whole successive crisis have taken place over the 1990s (with a reduction in alcohol prices, a drastic reduction in the production of alcohol cars, the lack of a policy for the sector), leading to the cooling of the investment capacity of the sector. We consider this a factor that favored the adoption of technological alternatives for the less innovative destination of vinasse.

The elements of the macroeconomic environment that influence this selection can be identified with low oil prices that have prevailed in those years. Low oil prices in the second half of 1980s reduced the competitiveness not only of alcohol but also made it difficult to expect economic pay-off from vinasse's biogas.

Recently, brand price volatility proceeds as instable macroeconomic signs adding more uncertainty to the private investment horizons. Investment in biogas recovery from vinasse treatment, as mentioned above, is of great importance for economic viability of anaerobic digestion. In this article, we design a whole set of analytical categories of vinasse disposal problem that should help to alert the reader that the conformation of a technological

trajectory is not in any way, single-determined. Oil prices, in this sense, represent only one aspect that contributes to this multi-determined phenomenon. As we see, persistently low prices of oil (and, in the case, especially diesel - also explained by the existence of fuel subsidies in the country - which is one element of a "political legal-institutional" nature) make higher the opportunity costs of investments in alternative energy technologies (such as biogas).

Elements of a political-legal and institutional environment that characterize the relevant selection environment for an alternative choice to fertigation include regulation on vinasse disposal and weak institutionalization of biotechnological research.

From the standpoint of the rules on vinasse disposal, as can be seen in Table 1, from the 78/79 crop it was forbidden to dump waste into surface waters, non compliance being punished with a fine.

The relevance of this legal ban seems clear. Margulis (1982) considers that fertigation has become an effectively diffused practice for biofuel industry by the time of the 1979/80 harvest, immediately after the release of the ban of vinasse disposal into surface waters.

Table 1 – Trend in regulation affecting vinasse disposal

Legislation	Description
Ordinance of the Ministry of Interior n° 323 29/11/1978	Ban vinasse disposal (direct and indirectly) into surface waters. Require distilleries to present projects for the implementation of systems to vinasse treatment or use.
Ordinance of the Ministry of Interior n° 158 November, 1980	Broadens the former Ordinance to the rest of the residual waters e liquid effluents from distilleries, and establishes deadlines and legal procedures for compliance projects elaboration and implementation. Restricts the access to government funding and incentives to the existence and approval of such projects.
CONAMA Resolution n° 0002 05/06/1984	Make it compulsory the presentation of studies and of a project with the solution to control the pollution caused by the effluents from distilleries and by sugarcane wash waters.
CONAMA Resolution n° 0001 23/01/1986	Make it compulsory the Environmental Impact Assessment and its respective Report for new distilleries and for any expansion projects for the existent ones.
Act n° 6.134 02/06/1988, art. 5°, Sao Paulo State Government	Establishes that liquid effluents, solid wastes and gases from any agro-industrial, industrial, commercial or other in nature, only can be disposed in forms that do not pose risks to groundwater.

Source: authors' elaboration.

Hassuda (1989) points out that the State of Sao Paulo counts with a legislation on the aquifer contamination since 1988 and that, so far, there were not any acts in the Federal level about the theme. Actions in this domain only were taken in 1999.

Rebouças (2002) calls attention to the fact that the project of the federal act to the regulation and management of groundwater (n° 7.127/86), had been under negotiation in the Parliament for over a decade, its promulgation being necessary to the regulation of Act 9.433 which involves water resources in general and promotes practical aspects for groundwater management.

Adopting an integrated policy to water resources (surface and groundwater) is a compromise that Brazil assumed in Agenda 21.

With regard to biotechnology, the initiative of major significance for the institutionalization of research in this area corresponds to the National Biotechnology (PRONAB) of the Department of Planning / National Research Council (SEPLAN / CNPq), in 1982.

Salles-Filho *et al* (1985) comment on the PRONAB:

"Despite the policy initiative that was launched, little was made of what was intended, since the unfavorable political and economic determinants that severely diminished the resources for Science and Technology development in the country."

PRONAB intended, through its extension in the State of Sao Paulo - the State Program for Biotechnology, to implement a series of actions to the development of anaerobic digestion (Anaerobic Digestion Sub-program). Among its short-term priorities, the Anaerobic Digestion Sub-program focused on basic and technological studies on anaerobic digestion of sewage from the sugar and alcohol mills - as well as urban waste and sewage. To cope with these aims, the Sub-program would have funded projects aiming at the integrated knowledge of microbiology and process biochemistry, anaerobic digesters optimization - more efficient from a technical-economic standpoint, granulation study in upflow digesters, evaluation of

building materials, consideration of the application of biogas and biofertilizer, spur of projects of deployment and diffusion of the developed technologies. This sub-program was not effectively implemented.

As previously mentioned, the existence of subsidies for diesel also constitutes a political and legal element to be considered, for it helps to boost the opportunity costs of alternative energy investments.

As regards to the elements of the natural environment, it is possible to say that just as the existence of watercourses in the neighborhood of plants and distilleries has long favored the practice of vinasse disposal into these waters, the availability of abundant land has also favored its view as a “natural” disposal for vinasse.

As elements of a social problem involved in the vinasse disposal, Plaza-Pinto (1999) refers to episodes of disputes involving plant owners and the public because of the "grout distillery" by the formers into the surface waters, with serious problems of local pollution. In his work of 1984, Rezende (*apud* Plaza-Pinto, 1999) reports ten major cases of contamination with vinasse published in the national press between 1943 and 1984. All episodes reported local catastrophes, such as large fish kills, water shortages, excessive proliferation of insects, public health problems and disruption of local economies dependent on fishing. One of the cases occurred in 1984, between the dams of Sobradinho and Moxotó in San Francisco River, was caused by the rupture of a stillage dam resulting in the spill of 45 billion m³ of vinasse in Tourão Creek, a tributary of San Francisco by Agrovale company. The result became known as the largest ecological disaster in the San Francisco River, killing 300 tonnes of fish. Although Agrovale tried to conceal the evidence of contamination, turning the land contaminated the river with tractors, it failed to dissuade the public and the authorities of its responsibility. According to author's account, the revolt of the river dwellers, whose survival depended on fishing, gave prominence to the episode. The option for fertigation, which

reduces the risk of contamination of water supplies in the short term, certainly has been helped by episodes like this, involving accidents caused by disruption of vinasse tanks leading to spillage of waste into surface waters.

5. Final discussion: understanding lock-in by multi-determined events

This paper presents the main results of a research focused on the analysis of the multiple factors involved in the formation of a technological trajectory for the solution of the vinasse destination problem in Brazil in the period of the 1970s to the 1990s.

We share a broader evolutionary view that cleaner technologies may be hold back by obstacles that must be understood by considering, above and beyond the essentially economic or technical factors – the top-down and the bottom-up approaches identified by Carrillo-Hermosilla *et al* (2009), the co-evolution of the change processes, both technological and institutional. Besides, along with Llerena & Matt (1999), Mulder & Van Den Bergh (2001) and Frenken *et al* (2004), we embraces the notion that it is essential to focus on path-dependent technological change within a context of co-evolving environmental, social and economic processes characterized both by irreversibility and uncertainty, rather than on maximization and equilibrium, emphasized by mainstream economics.

This approach must prevent us to say that the ultimate reason that explains why fertigation came to be the "winning" solution to the vinasse disposal problem could be its cost-effectiveness. This result could be achieved by means of a conventional cost-benefit analysis. Studying the economic feasibility of alternatives at the time, Margulis (1982) was enthusiastic about fertigation, but did not rule out the possibility that, over time, anaerobic digestion – still immature from technical point of view – could came to be an economically viable option.

After almost thirty years from the Margulis' evaluation, fertigation remains the most widespread solution in the country, and have been consolidated in practice as the "solution" to the vinasse disposal problem.

Van Lente (1997) found that biotechnology had gained widespread attention by the promise of helping to reduce pollution and to the energy production, opening up new perspectives to contribute to sustainable development and the process of the greening of industry. These possibilities could characterize the technological opportunities, as refereed by Montalvo (2007), and could also be understood as a business opportunity. During two decades, comprising the years of 70s and 80s, anaerobic digestion brightly represented a possibility for this promise to be fulfilled in Brazil.

It is clear that the promises of biotechnology were not restricted to Brazil or to a specific industry. The widespread belief in the emergence of a new paradigm that could replace the "old" chemical pathways for the production of pharmaceuticals, agrochemicals and a wide range of new products impacting across major industrial sectors were supported, as indicated by Van Lente (1997) by undeniable evidence: significant scientific developments, the reduction of the gap between basic and applied science, the climb of incentives for equipment development, the magnitude of inter-sectoral impacts forecasted. These would be evidence of an overly optimistic prognosis with respect to the development of biotechnology and its applications, as in the case of anaerobic digestion?

By that time (late 1990s) and in the case of the vinasse biodigestion, the gap between the promise of biotechnology and their actual achievements is directly proportional to the general disappointment after the euphoria experienced in the early 80s.

In our particular case study, we list a number of elements that help us not to confine our understanding of the upholding of fertigation as the resolution to vinasse problem to a too simplistic cost-benefit explanation. In the paragraphs below, we point out elements that

contribute to the continuity of the technological trajectory of fertigation and the exclusion of alternative anaerobic digestion and advance a few comments on the possibility of breaking the consolidated trajectory.

Initially, it would be logical to assume that a cost-benefit analysis, even if not formalized by the users, had been largely in favor of fertigation, along with its high cost-efficiency. The work of Margulis (1982) is illuminating on these points.

As it was already pointed out, the sugarcane agro-industry, with traditionally simple technology, is not characterized by a significant technological dynamics: let us remember, once more, the proportion of generation vinasse and ethanol remained practically unchanged since 1975 despite the investments offered by ProAlcool. Probably the manufacture side of the sector could be best described, by the time, by organizational rigidity and resistance to innovation, which is mostly dependent of suppliers.

The same is not true, nevertheless, when it comes to the innovative efforts which are traditionally by a web of mainly, but not exclusively, public supported agents related to the sugarcane industry in the domain of genetic improvement of the plant, combating pests, agricultural and harvesting techniques, impact of the cultivation on the environment, and, more recently, technologies concerning the manufacture of ethanol, including hydrolysis and fermentation (FAPESP, 2007).

The urgency to give a destination (an alternative to surface water) to huge amounts of vinasse within extremely tight deadlines - posted by Ordinance of the Ministry of Interior in 1978-79 – appears to have benefited fertigation; it can be said that this was the most accommodating alternative to the selection environment at the time.

The developments experienced by fertigation through the 70s and 80s provided the technological trajectory the exploitation of economies of scale, in the way discussed by many authors (David, 1985; Arthur, 1988), but also of the economies of scope. In terms of scale, the

development of dispensing systems has expanded the acreage that could receive the waste in an economically efficient way. In terms of scope, the study of rates of disposal has made possible the substitution of the waste for chemical inputs (fertilizer).

We also believe that one can talk, to a certain degree, of technological interdependence (between fertigation itself and the application technology and infrastructure) to the extent that widespread use of fertigation just followed the ban on the release of the waste into surface waters, i.e. from the 1979/80 harvest. This possibility is reinforced by the need for closer links between users (producers of sugarcane) and suppliers of fertigation equipment. This observation is consistent with the notion of system integration of the technological solution put forward by authors like Kemp *et al* (1998). The narrowing of these links tends to favor the qualification of users in the operation of such equipment and the establishment of a certain infrastructure (canals or pipelines, investments in trucks, tanks and pumps etc.), which contributes to strengthen the advancement of the technological trajectory, as was already pointed out by David (1985) and Arthur (1988).

The achievement of these investments, on the other hand, implies the emergence of sunk costs. Although it is difficult to specify precisely the dimension of these costs, one can say that their importance tends to be greater when considering the crisis faced by the sector at the time, characterized by the stability of the selection environment (with diminishing demand for ethanol as fuel for vehicles and without further pressures for technological change in this area).

The spread of fertigation should also have caused the emergence of some positive network externalities and this observation is in accordance with David (1985) and Arthur (1988). That is to say, the adoption of the application of natural vinasse in soil for the first producers may have performed as a "demonstration effect", influencing other producers to adopt it as well.

One could even speak of a lack of information on alternative technologies or of a lack of knowledge about how to use this information, as it is been considered by Carrillo-Hermosilla *et al* (2009).

In any case, a technology in its initial stages of development, as was the case of anaerobic digestion, requires an extremely low degree of risk aversion from early adopters. Additionally, in the specific case of this technology, it should require a very great believe in the promising biotechnology paradigm. Indeed, it would be surprising a surge on its adoption in those years. These conditions, namely the low degree of risk aversion and the great believe in biotechnology, seemed to be found only in a few plants in Brazil at the time: PAISA , São Martinho and Boa Vista.

As we saw in the case of development of anaerobic digestion, it is not true that technological opportunities have been opened by new scientific knowledge. It was necessary for researchers to engage to answer questions more properly scientific, while seeking technical solutions to the trade-offs faced.

The expectations on technology change, or on the ability to achieve the promises of biotechnology, however, were strong enough to justify these efforts or support of the bets.

One might think, still, the possibility of coexistence between the anaerobic digestion of vinasse and the fertigation. Perhaps both could, to some extent, be considered complementary rather than properly competing.

The adoption by a small number of plants can be considered as indicative of the possibility of creating certain "market niches" in which penetration of innovation meet less resistance. The use of biogas for dry yeast can be understood as an attempt to exploit a market niche.

The efforts of anaerobic digestion told, in this sense, the willingness of established plants (PAISA, São Martinho and Boa Vista) to engage in innovation and proactive strategy

to internalize the treatment of vinasse with biogas production. But we also see the entry of at least one new firm, Biometano, as a supplier of technological capabilities.

Those proved not be sufficient conditions to break the pattern of technological solution that was already spreading (the fertigation).

The main technological bottlenecks faced by fertigation were being overcome in a fairly efficient way. One could not identify any new significant problem area challenging its advance, so that, after having surpassed some technical difficulties, not only fertigation has not faced any economic restrictions but, moreover, has provided a simple, cheap and profitable way to discard huge quantities of the polluting waste. A real Columbus' egg.

The changes that have taken place in the selection environment proved, on one hand, to be favorable to the fertigation trajectory (through the tight deadlines for the compliance with new environmental regulation) and, on the other hand, hostile to anaerobic digestion (due to the weak institutionalization of biotechnology research in the country).

For many, the problem of disposal of the waste is currently considered resolved. It is not necessary here to retrieve all the advantages attributed to fertigation.

It is undeniable that fertigation has constituted an extremely effective response when compared to vinasse release into rivers. The effects of this practice were sufficiently obvious not to raise scientific disputes and to justify the tightness of deadlines to compliance.

However, that certainty with respect to environmental impacts of the release of vinasse into surface waters is unlikely to be repeated, as we have already pointed out, in the case of the potential environmental effects of its disposal on the ground.

We saw that the alcohol agro-industry activities have been identified as one of the vulnerability factors of São Paulo aquifers and some studies have called attention to the risk of groundwater contamination by vinasse disposal in soil. Soil salinisation is also identified as a danger associated with this practice.

Both the groundwater contamination and soil salinisation are treated as "risks" and the studies on the matter are still inconclusive. Given these risks, there remains the question if the fertigation would not only be a temporary, short-term solution given the emergency of the vinasse disposal problem. It seems, likewise, that it was premature to abandon efforts to develop the anaerobic digestion of the waste.

What is clear is that more research was needed to clarify these questions. In the absence of such a clarification, we are left with the impression that we are dealing with a case where the solution retained is a quick and inexpensive way to get rid of the sugarcane agro-industry annoying and perhaps even dangerous waste. Worse than that, not to investigate these issues implies leaving to future generations the task of discovering whether the infiltration of vinasse in soil actually contributes to soil salinisation and groundwater contamination.

6. References

- ARTHUR, W. B. (1988) Competing technologies: an overview. In: Dosi, G. *et al Technical change and economic theory*. London: Pinter Publ.
- CARRILLO-HERMOSILLA, J.; GONZÁLEZ, P. D. R.; KÖNNÖLÄ, T. (2009) *Eco-innovation: when sustainability and competitiveness shake hands*. New York: Palgrave Macmillan.
- CORAZZA, R. I. Políticas públicas para tecnologias mais limpas: uma análise das contribuições da economia do meio ambiente. 2001. Tese (Doutorado em Política Científica e Tecnológica) no Departamento de Política Científica e Tecnológica, Instituto de Geociências, Universidade de Campinas, Campinas, 2001.

- CORTEZ, L. A. B.; LARSON, D. L. Environmental aspects of the alcohol program in Brazil. Paper presented at the *ASAE Annual International Meeting*, July 11-16, 1998, Orlando, Florida, USA.
- COWAN, R.; HULTÉN, S. (1996) Escaping Lock-In: The Case of the Electric Vehicle, *Technological Forecasting and Social Change* (53): 61-79.
- COWAN, R.; KLINE, D. (1996) The Implications of Potential 'Lock-In' in Markets for Renewable Energy. *National Renewable Energy Laboratory*, Golden, CO.
- CRUZ, R. L. *Efeito da aplicação de vinhaça sobre o solo e água subterrânea*. Dissertação (Mestrado em Hidráulica e Saneamento) na Escola de Engenharia de São Carlos, Universidade de São Paulo, 1991.
- DAVID, P. (1985) Clio and the economics of QWERTY. *American Economic Review AEA Papers and Proceedings*, (75):332-337.
- DOLFSMA, W.; LEYDESDORFF, L. (2009) Lock-in and Break-out from Technological Trajectories: modeling and policy implications, *Technological Forecasting and Social Change*, 76(7): 932-941.
- DOSI, G. (1982) Technological paradigms and technological trajectories: a suggested interpretation of the determinants and directions of technical change. *Research Policy* (11):147-162.
- FAPESP (The State of São Paulo Research Foundation) (2007) Brazil: world leader in sugarcane and ethanol knowledge and technology – Fapesp's contributions. São Paulo: Fapesp. Available in: http://www.fapesp.br/publicacoes/livro_etanol_ing.pdf
- FOXON, T.; GROSSA, R.; CHASEB, A.; HOWESB, J.; ARNALLC, A.; ANDERSON, S. (2005) UK innovation systems for new and renewable energy technologies: drivers, barriers and systems failures, *Energy Policy* (33): 2123–2137.

- FRENKEN, K., HEKKERT, M., GODFROIJ, P. (2004) R&D portfolios in environmental friendly automotive propulsion: variety, competition and policy implications. *Technological Forecasting & Social Change* 71 (5), 485– 507.
- GLOEDEN, E.; CUNHA, R.C.A.; FRACCAROLLI, M.J.B.; CLEARY, R.W. The behaviour of vinasse constituents in the unsaturated zones in the Botucatu aquifer recharge area. *Water Science and Technology* 24(11):147-157. 1991.
- GUARNIERI, L. C.; JANUZZI, G. de M. Proálcool: impactos ambientais. *Revista Brasileira de Energia* 2(2):147-161, 1992.
- HASSUDA, S. *Impactos da infiltração da vinhaça de cana no aquífero de Bauru*. Dissertação (Mestrado em Geociências) Universidade de São Paulo, 1989.
- HIRATA, R. C. A.; BASTOS, C.; ROCHA, G.; GOMES, D.; IRITANI, M. Groundwater pollution risk and vulnerability map of the state of São Paulo, Brazil. *Water Science and Technology* 24(11):159-169, 1991.
- HOOGMA, R.; KEMP, R.; SCHOT, J.; TRUFFER, B. (2002) *Experimenting for Sustainable Transport*. London: Spon Press.
- JACOBSON, S.; JOHNSON, A. (2000) The diffusion of renewable energy technology: an analytical framework and key issues for research. *Energy Policy*, 28:625-640.
- JACOBSSON, S.; BERGEK, A. (2004) Transforming the energy sector: the evolution of technological systems in renewable energy technology. *Industrial and Corporate Change*, 13 (5): 815-849.
- KEMP, R., SCHOT, J.; HOOGMA, R. (1998) Regime shifts to sustainability through processes of niche formation: the approach of strategic niche management. *Technology Analysis and Strategic Management*, 10(2):175–197.
- LA ROVERE, E. L. Les impacts sociaux et écologiques du Plan Alcool Brésilien. *Économie et humanisme* (260): 36-47, 1981.

- LLERENA, P., MATT, M. (1999) Inter-organizational collaborations: the theories and their policy implications. In: Gambardella, A., Malerba, F. (Eds.), *The Organization of Economic Innovation in Europe*. Cambridge University Press, pp. 179–201.
- MARGULIS, S. Vinhoto: poluição hídrica, perspectivas de aproveitamento e interação com o modelo matemático de biomassa. *Texto para discussão de energia n. 10*. Rio de Janeiro: IPEA/INPES, 1982.
- MONTALVO, C. (2007) General wisdom concerning the factors affecting the adoption of cleaner technologies: a survey 1990-2007, *Journal of Cleaner Production*, 16(1):7-13.
- MULDER, P., VAN DEN BERGH, J. (2001) Evolutionary economic theories of sustainable development. *Growth and Change* (32): 110– 134.
- PLAZA PINTO, C. (1999) *Tecnologia da digestão anaeróbia da vinhaça e desenvolvimento sustentável*. Dissertação (Mestrado em Engenharia Mecânica). Campinas: FEM/Unicamp.
- RAVEN, R. P. J. M. (2005) *Strategic Niche Management for Biomass* A comparative study on the experimental introduction of bioenergy technologies in the Netherlands and Denmark. Eindhoven: Technische Universiteit Eindhoven.
- REBOUÇAS, A. C. A política nacional de recursos hídricos e as águas subterrâneas. *Rev. Águas Subterrâneas*, v. 16, 2002.
- RIGHETTO, A. M.; CRUZ, R. L.; NOGUEIRA, M. A. Experimental investigation of soil and groundwater impacts caused by vinasse disposal. *Water Science Technology* 24(11):77-85, 1991.
- SALLES-FILHO, S. L. M.; CERANTOLA; W. A.; ALVARES, V. M.P. As promessas da biotecnologia para a América Latina: um alerta. *Revista Brasileira de Tecnologia* 16(6):13-27, 1985

- SCHELEDER, E. M. M. *A questão do álcool combustível*. Seminário sobre a indústria canavieira, Campinas: Unicamp/IE, 1998. http://www.eco.unicamp.br/pesquisa/frame_inic.html. (consultado em 03/1999).
- SZMRECSÁNYI, T. Tecnologia e degradação ambiental: o caso da agroindústria canavieira no Estado de São Paulo. *Informações Econômicas* 24(10):73-81, 1994.
- UNRUH, G. C. (2000) Understanding carbon lock-in. *Energy Policy*, 28:817-830.
- VAN DEN BERGH, J. C. J. M.; FABER, A.; IDENBURG, A.; OOSTERHUIS, F. H. (2006) Survival of the greenest: evolutionary economics and policies for energy innovation, *Environmental Sciences*, 3(1): 57 – 71.
- VAN LENTE, H. *Towards an integrated conceptual framework: modern biotechnology and the greening of industry*. Prepared for the European Commission. Seville, Spain: JRC Institute for Prospective Technological Studies, 1997.
- VERBONG, G.; GEELS, F. (2006) The ongoing energy transition: Lessons from a socio-technical, multi-level analysis of the Dutch electricity system (1960–2004), *Energy Policy* (35): 1025–1037.